## **DRAFT** February 2008

## **Calculation of Potential Risk from Consumption of Breast Milk**

Chemical	Cf (mg/kg)	SFo (mg/kg/day)-1	RfD (mg/kg/day)	h (days)	ADDm (mg/kg/day)	Cmf (mg/kg-lipid	ADDca-c d) (mg/kg/day)	ADDnc-c (mg/kg/day)	Mother ELCRm	Child ELCRc	Mother HQm	Child HQc
PCBs	1.6	2	0.00002	2555	0.0034	38	0.00144	0.101	3.0E-03	2.9E-03	172	5030
							Factor above ac	ceptable level =	2951	2874	172	5030
PCBs TEQ	1.90E-05	1.50E+05	NA	2555	4.1E-08	4.5E-04	1.7E-08	1.2E-06	2.6E-03	2.6E-03		
							Factor above ac	ceptable level =	2628	2560		
Dioxin TEQ	5.10E-06	1.50E+05	NA	2555	1.1E-08	1.2E-04	4.6E-09	3.2E-07	7.1E-04	6.9E-04		
DIOXIII I EQ	0.102 00	1.002100	14/1	2000	1.12 00	_		ceptable level =		687		
DDT	0.07	0.04	0.0005	400	0.0000	0.00	0.05.00	0.45.04	0.05.05	4.05.00	0.00	0.44
DDT	0.07	0.34	0.0005	120	0.0002	0.08	3.0E-06 Factor above ac	2.1E-04 ceptable level =	2.2E-05 22	1.0E-06 1.0	0.30 0.30	0.41 0.41
							i actor above ac	coptable level =		1.0	0.00	0.41

## Notes:

Site-wide, whole-body 95UCL concentration in small mouth bass (Cf) taken from Round 2 Report, Appendix F, Table 3-14.

Acceptable levels are ELCR = 1E-6 and HQ = 1

ELCRm adjusted to 30-year exposure

## **Equations**

Atc

Atnc

$$\begin{split} & ADDm = (Cf \ x \ Irf \ x \ Conv \ x \ Ff) \ / \ BWm \\ & ADDca-c = (Cmf \ x \ Irm \ x \ f3 \ x \ f4 \ x \ Edc \ x \ Efc) \ / \ (Atc \ x \ BWc) \end{split}$$

25550 days

365 days

ELCRm = ADDm x Sfo ELCRc = ADDca-c x Sfo Cmf = (ADDm x h x f1) / (In2 x f2)

ADDnc-c =  $(Cmf \times Irm \times f3 \times f4 \times Edc \times Efc) / (Atnc \times BWc)$ 

HQm = ADDm / RfD HQc = ADDnc-c / RfD

Averaging time - carcinogens (70 years)

Averaging time - noncarcinogens (Edc x Efc)

Default V	alues		Calculated	Calculated Values					
Cf	chemical specific mg/kg	Concentration of chemical in fish	ADDm	mg/kg/day	Average Daily Dose to mother				
Irf	142 mg/day	Mother's ingestion rate of fish	Cmf	mg/kg-lipid	Chemical concentration in milkfat				
Conv	0.001 kg/mg	Conversion factor	ADDca-c	mg/kg/day	Average Daily Dose to breast-feeding child, cancer				
Ff	1 fraction	Fraction of fish contaminated	ADDnc-c	mg/kg/day	Average Daily Dose to breast-feeding child, non-cancer				
BWm	66 kg	Body weight of mother	ELCRm	risk	Excess Lifetime Cancer Risk to mother				
h	chemical specific days	Half-life of chemical in body	ELCRc	risk	Excess Lifetime Cancer Risk to child				
Fone	0.9 fraction	Fraction of ingested chemical stored in fat	HQm	quotient	Hazard Quotient to mother				
Ftwo	0.3 fraction	Fraction of mother's weight that is fat	HQc	quotient	Hazard Quotient to child				
Irm	0.69 kg/day	Infant's ingestion rate of milk							
Fthree	0.04 fraction	Fraction of breast milk that is fat							
Ffour	0.9 fraction	Fraction of ingested chemical that is absorbed							
Edc	1 year	Exposure duration of breast-feeding child							
Efc	365 days/year	Exposure frequency of breast-feeding child							

BWc 9.4 kg Body weight of child Sfo chemical specific (mg/kg/day)-1 Slope Factor - oral RfD chemical specific mg/kg/day Reference Dose - oral